

# A CHARACTERIZATION OF SUBSPACES AND QUOTIENTS OF REFLEXIVE BANACH SPACES WITH UNCONDITIONAL BASES

---

W. B. JOHNSON and BENTUO ZHENG

## Abstract

*We prove that the dual or any quotient of a separable reflexive Banach space with the unconditional tree property (UTP) has the UTP. This is used to prove that a separable reflexive Banach space with the UTP embeds into a reflexive Banach space with an unconditional basis. This solves several longstanding open problems. In particular, it yields that a quotient of a reflexive Banach space with an unconditional finite-dimensional decomposition (UFDD) embeds into a reflexive Banach space with an unconditional basis.*

## 1. Introduction

It has long been known that Banach spaces with unconditional bases as well as their subspaces are much better behaved than general Banach spaces and that many of the reflexive spaces (including  $L_p(0, 1)$ ,  $1 < p < \infty$ ) which arise naturally in analysis have unconditional bases. It is, however, difficult to determine whether a given Banach space has an unconditional basis or if it embeds into a space that has an unconditional basis. Two problems, considered important since at least the 1970s, stand out.

- (a) Give an intrinsic condition on a Banach space  $X$  which is equivalent to the embeddability of  $X$  into a space with an unconditional basis.
- (b) Does every complemented subspace of a space with an unconditional basis have an unconditional basis?

Problem (b) remains open, but in this article, we provide a solution to problem (a) for reflexive Banach spaces. This characterization also yields that a quotient of a reflexive space with an unconditional basis embeds into a reflexive space with unconditional basis, which solves another problem from the 1970s. Here, some condition on the space with an unconditional basis is needed because every separable Banach space is a quotient of  $\ell_1$ .

DUKE MATHEMATICAL JOURNAL

Vol. 141, No. 3, © 2008 DOI 10.1215/00127094-2007-003

Received 12 December 2006. Revision received 16 May 2007.

2000 *Mathematics Subject Classification*. Primary 46B03; Secondary 46B20.

Johnson's work partially supported by National Science Foundation grants DMS-0200690 and DMS-0503688.

Zheng's work partially supported by National Science Foundation grants DMS-0200690 and DMS-0503688.

There is, of course, quite a lot known concerning problems (a) and (b). For example, Pełczyński and Wojtaszczyk [15, Theorem 1.1] proved that if  $X$  has an unconditional expansion of identity (i.e., a sequence  $(T_n)$  of finite-rank operators such that  $\sum T_n$  converges unconditionally in the strong operator topology to the identity on  $X$ ), then  $X$  is isomorphic to a complemented subspace of a space that has an unconditional finite-dimensional decomposition (UFDD). Later, Lindenstrauss and Tzafriri [11, Theorem 1.g.5] showed that every space with a UFDD embeds (not necessarily complementably) into a space with an unconditional basis. As regards reflexive spaces, it was shown in [4, Theorem 3.1] using a result from [1, Lemma 1] (and answering a question from that article), that if a reflexive Banach space embeds into a space with an unconditional basis, then it embeds into a reflexive space with an unconditional basis. As regards the quotient problem mentioned above, Feder [3, Theorem 4] gave a partial solution by proving that if  $X$  is a quotient of a reflexive space that has a UFDD and  $X$  has the approximation property, then  $X$  embeds into a space with an unconditional basis.

It is well known and easy to see that if a Banach space  $X$  embeds into a space with an unconditional basis, then  $X$  has the unconditional subsequence property; that is, there exists a  $K > 0$  such that every normalized weakly null sequence in  $X$  has a subsequence that is  $K$ -unconditional. In fact, failure of the unconditional subsequence property is the only known criterion for proving that a given reflexive space does not embed into a space with an unconditional basis. However, in Section 3, we construct a Banach space that has the unconditional subsequence property but does not embed into a Banach space that has an unconditional basis. This is not surprising, given previous examples of Odell and Schlumprecht [12]. Moreover, Odell and Schlumprecht have taught us that when a subsequence property is replaced with the corresponding “branch of a tree” property (see [12, introduction]), the result is a stronger property that sometimes can be used to give a characterization of spaces that embed into a space with some kind of structure. The relevant property for us here is the unconditional tree property (UTP), and Odell and Schlumprecht’s beautiful results are essential tools for us in applying it.

We use standard Banach space theory terminology, such as can be found in [11].

## 2. Main results

### *Definition 2.1*

Let  $[\mathbf{N}]^{<\omega}$  denote all finite subsets of the positive integers. By a normalized weakly null tree, we mean a family  $(x_A)_{A \in [\mathbf{N}]^{<\omega}} \subset S_X$  with the property that every sequence  $(x_{A \cup \{n\}})_{n \in \mathbf{N}}$  is weakly null. Let  $A = \{n_1, \dots, n_m\}$  with  $n_1 < \dots < n_m$ , and let  $B = \{j_1, \dots, j_r\}$  with  $j_1 < \dots < j_r$ . Then we say that  $A$  is an initial segment of  $B$  if  $m \leq r$  and  $n_i = j_i$  when  $1 \leq i \leq m$ . The tree order on  $(x_A)_{A \in [\mathbf{N}]^{<\omega}}$  is given by  $x_A \leq x_B$  if  $A$  is an initial segment of  $B$ . A branch of a tree is a maximal linearly

ordered subset of the tree under the tree order. We say that  $X$  has the  $C$ -UTP if every normalized weakly null tree in  $X$  has a  $C$ -unconditional branch for some  $C > 0$  and that  $X$  has the UTP if  $X$  has the  $C$ -UTP for some  $C > 0$ .

*Remark 2.2*

Odell, Schlumprecht, and Zsák proved in [14, Proposition 2.2] that if every normalized weakly null tree in  $X$  admits a branch that is unconditional, then  $X$  has the  $C$ -UTP for some  $C > 0$ . A simpler proof appears in the preprint of Haydon, Odell, and Schlumprecht [5]. There is, therefore, no ambiguity when using the term UTP.

Given a finite-dimensional decomposition (FDD)  $(E_n)$ ,  $(x_n)$  is said to be a *block sequence* with respect to  $(E_n)$  if there exists a sequence of integers  $0 = m_1 < m_2 < m_3 < \dots$  such that  $x_n \in \bigoplus_{j=m_n}^{m_{n+1}-1} E_j, \forall n \in \mathbf{N}$ . Further,  $(x_n)$  is said to be a *skipped block sequence* with respect to  $(E_n)$  if there exists a sequence of increasing integers  $0 = m_1 < m_2 < m_3 < \dots$  such that  $m_n + 1 < m_{n+1}$  and  $x_n \in \bigoplus_{j=m_n+1}^{m_{n+1}-1} E_j, \forall n \in \mathbf{N}$ . Let  $\delta = (\delta_i)$  be a sequence of positive numbers decreasing to zero. We say that  $(y_n)$  is a  $\delta$ -*skipped block sequence* with respect to  $(E_n)$  if there is a skipped block sequence  $(x_n)$  such that  $\|y_n - x_n\| < \delta_n \|y_n\|, \forall n \in \mathbf{N}$ . We say that  $(F_n)$  is a *blocking* of  $(E_n)$  if there is a sequence of increasing integers  $0 = k_0 < k_1 < \dots$  such that  $F_n = \bigoplus_{j=k_{n-1}+1}^{k_n} E_j$ .

*Definition 2.3*

Let  $X$  be a Banach space with an FDD  $(E_n)$ . If there exists a  $C > 0$  such that every skipped block sequence with respect to  $(E_n)$  is  $C$ -unconditional, then we say that  $(E_n)$  is an *unconditional skipped block FDD (USB FDD)*.

The following is a blocking lemma of Johnson and Zippin (see [10] or [11, Proposition 1.g.4(a)]) which is used later.

LEMMA 2.4

Let  $T : X \rightarrow Y$  be a bounded linear operator. Let  $(B_n)$  be a shrinking FDD of  $X$ , and let  $(C_n)$  be an FDD of  $Y$ . Let  $(\delta_n)$  be a sequence of positive numbers tending to zero. Then there are blockings  $(B'_n)$  of  $(B_n)$  and  $(C'_n)$  of  $(C_n)$  such that for every  $x \in B'_n$ , there is a  $y \in C'_{n-1} \oplus C'_n$  such that  $\|Tx - y\| \leq \delta_n \|x\|$ .

The lemma above actually works for any further blockings of  $(B'_n)$  and  $(C'_n)$ . To be more precise, we have the following stronger result, which is actually a formal consequence of Lemma 2.4 as stated.

LEMMA 2.5

Let  $T : X \rightarrow Y$  be a bounded linear operator. Let  $(B_n)$  be a shrinking FDD of  $X$ , and let  $(C_n)$  be an FDD of  $Y$ . Let  $(\delta_n)$  be a sequence of positive numbers tending

to zero. Then there are blockings  $(B'_n)$  of  $(B_n)$  and  $(C'_n)$  of  $(C_n)$  such that for any further blockings  $(\tilde{B}_n)$  of  $(B'_n)$  with  $\tilde{B}_n = \bigoplus_{i=k_n}^{k_{n+1}-1} B'_i$  and  $(\tilde{C}_n)$  of  $(C'_n)$  with  $\tilde{C}_n = \bigoplus_{i=k_n}^{k_{n+1}-1} C'_i$  and for any  $x \in \tilde{B}_n$ , there is a  $y \in \tilde{C}_{n-1} \oplus \tilde{C}_n$  such that  $\|Tx - y\| \leq \delta_n \|x\|$ .

*Proof*

Let  $(\delta_i)$  be a sequence of positive numbers decreasing to zero. Let  $(\tilde{\delta}_i)$  be another sequence of positive numbers that go to zero so fast that  $\sum_{j=i}^\infty \tilde{\delta}_j < \delta_i/2\lambda$ , where  $\lambda$  is the basis constant for  $(B_n)$ . By Lemma 2.4, we get blockings  $(B'_n)$  of  $(B_n)$  and  $(C'_n)$  of  $(C_n)$ , so that for every  $x \in B'_n$ , there is a  $y \in C'_{n-1} \oplus C'_n$  such that  $\|Tx - y\| \leq \tilde{\delta}_n \|x\|$ . Let  $\tilde{B}_n = \bigoplus_{i=k_n}^{k_{n+1}-1} B'_i$  and  $\tilde{C}_n = \bigoplus_{i=k_n}^{k_{n+1}-1} C'_i$  be blockings of  $(B'_n)$  and  $(C'_n)$ . Let  $x \in \tilde{B}_n$ . Then we can write  $x = \sum_{i=k_n}^{k_{n+1}-1} x_i, x_i \in B'_i$ . So by our construction of  $(B'_n)$  and  $(C'_n)$ , there are  $y_i \in C'_{i-1} \oplus C'_i, k_n \leq i \leq k_{n+1} - 1$ , such that  $\|Tx_i - y_i\| \leq \tilde{\delta}_i \|x_i\|, k_n \leq i \leq k_{n+1} - 1$ . Let  $y = \sum_{i=k_n}^{k_{n+1}-1} y_i \in \tilde{C}_{n-1} \oplus \tilde{C}_n$ . Then we have

$$\|Tx - y\| \leq \sum_{i=k_n}^{k_{n+1}-1} \tilde{\delta}_i \|x_i\| \leq \sum_{i=k_n}^{k_{n+1}-1} 2\lambda \tilde{\delta}_i \|x\| \leq \delta_n \|x\|. \quad \square$$

The following convenient reformulation of Lemma 2.4 is also used later (see [9] and [10] or [13]).

LEMMA 2.6

Let  $T : X \mapsto Y$  be a bounded linear operator. Let  $(B_n)$  be a shrinking FDD for  $X$ , and let  $(C_n)$  be an FDD for  $Y$ . Let  $(\delta_i)$  be a sequence of positive numbers decreasing to zero. Then there is a blocking  $(B'_n)$  of  $(B_n)$  and a blocking  $(C'_n)$  of  $(C_n)$  such that for any  $x \in B'_n$  and any  $m \neq n, n - 1$ ,

$$\|Q_m(Tx)\| \leq \delta_{\max\{m,n\}} \|x\|,$$

where  $Q_j$  is the canonical projection from  $Y$  onto  $C'_j$ .

*Remark 2.7*

The qualitative content of Lemma 2.6 is that there are blockings  $(B'_n)$  of  $(B_n)$  and  $(C'_n)$  of  $(C_n)$  such that  $T B'_n$  is essentially contained in  $C'_{n-1} + C'_n$ .

Our first theorem says that the UTP for reflexive Banach spaces passes to quotients. It plays a key role in this article and involves Lemmas 2.4–2.6 as well as results and ideas of Odell and Schlumprecht.

Let us explain the sketch of the proof of the special case where  $Y$  is a reflexive space with the UTP and  $Y$  has an FDD  $(E_n)$  while  $X$  is a quotient of  $Y$  which has an FDD  $(V_n)$ . Since  $Y$  has the UTP, by Odell and Schlumprecht’s fundamental result

([12, Theorem 3.3]) there is a blocking  $(F_n)$  of the  $(E_n)$  which is a USB FDD. Then we use the “killing the overlap” technique of [6] to get a further blocking  $(G_n)$  so that any norm 1 vector  $y$  is a small perturbation of the sum of a skipped block sequence  $(y_i)$  with respect to  $(F_n)$  and  $y_i \in G_{i-1} \oplus G_i$ . Let  $Q : Y \mapsto X$  be the quotient map. Using Lemma 2.5 and passing to a further blocking, without loss of generality, we assume that  $QG_i$  is essentially contained in  $H_{i-1} + H_i$ , where  $(H_i)$  is the corresponding blocking of  $(V_n)$ . Let  $(x_A)$  be a normalized weakly null tree in  $X$ . We then choose a branch  $(x_{A_i})$  so lacunary that  $(x_{A_i})$  is a small perturbation of a block sequence of  $(H_n)$ , and for each  $i$  there is at least one  $H_{k_i}$  between the essential support of  $x_{A_i}$  and  $x_{A_{i+1}}$ . Let  $x = \sum a_i x_{A_i}$  with  $\|x\| = 1$ . Considering a preimage  $y$  of  $x$  under the quotient  $Q$  from  $Y$  onto  $X$  (with  $\|y\| = 1$ ), by our construction we can essentially write  $y$  as the sum of  $(y_i)$ , where  $(y_i)$  is a skipped block sequence with respect to  $(F_n)$ . Since  $(F_n)$  is a USB,  $(y_i)$  is unconditional. By passing to a suitable blocking  $(z_i)$  of  $(y_i)$  and then using Lemma 2.5, it is not hard to show that  $Qz_i$  is essentially equal to  $a_i x_{A_i}$ . Noticing that  $(z_i)$  is unconditional, we conclude that  $(x_{A_i})$  is also unconditional.

For the general case where  $X$  and  $Y$  do not have an FDD, we have to embed them into some superspaces with FDD. The difficulty is that when we decompose a vector in  $Y$  as the sum of disjointly supported vectors in the superspace, we do not know that the summands are in  $Y$ . The same problem occurs for vectors in  $X$ . This makes the proof rather technical and requires many computations.

**THEOREM 2.8**

*Let  $X$  be a quotient of a separable reflexive Banach space  $Y$  with the UTP. Then  $X$  has the UTP.*

*Proof*

By Zippin’s result (see [17]),  $Y$  embeds isometrically into a reflexive space  $Z$  with an FDD. A key point in the proof is that Odell and Schlumprecht proved (see [13, Proposition 2.4]) that there is a further blocking  $(G_n)$  of the FDD for  $Z$ ,  $\delta = (\delta_i)$ , and a  $C > 0$  such that every  $\delta$ -skipped block sequence  $(y_i) \subset Y$  with respect to  $(G_i)$  is  $C$ -unconditional. Let  $\lambda$  be the basis constant for  $(G_n)$ .

Since  $X$  is separable, we can regard  $X$  as a subspace of  $L_\infty$ . Let  $\epsilon > 0$ . We may assume that

- (a)  $\sum_{j>i} \delta_j < \delta_i$ ;
- (b)  $i \delta_i < \delta_{i-1}$ ; and
- (c)  $\sum \delta_i < \epsilon$ .

Let  $Q$  be a quotient map from  $Y$  onto  $X$  (which can be extended to a norm 1 map from  $Z$  into  $L_\infty$ , and we still denote it by  $Q$ ). Further,  $QZ$ , as any separable subspace of  $L_\infty$ , is contained in some superspace isometric to  $C(\Delta)$  with monotone basis  $(v_i)$ . Here,  $\Delta$  is the Cantor set.

Let  $(x_A)$  be a normalized weakly null tree in  $X$ . Then we let  $(E_n)$  and  $(F_n)$  be blockings of  $(G_i)$  and  $(v_i)$ , respectively, which satisfy the conclusions of Lemmas 2.5 and 2.6. Using the “killing the overlap” technique (see [13, Proposition 2.6]), we can find a further blocking  $(\tilde{E}_n = \bigoplus_{i=l(n)+1}^{l(n+1)} E_i)$  so that for every  $y \in S_Y$ , there exist  $(y_i) \subset Y$  and integers  $(t_i)$  with  $l(i - 1) < t_i \leq l(i)$  for all  $i$  such that

- (I)  $y = \sum y_i$ ;
- (II) for  $i \in \mathbf{N}$ , either  $\|y_i\| < \delta_i$  or  $\|\sum_{j=t_{i-1}+1}^{t_i-1} P_j y_i - y_i\| < \delta_i \|y_i\|$ ;
- (III)  $\|\sum_{j=t_{i-1}+1}^{t_i-1} P_j y - y_i\| < \delta_i$ ; and
- (IV)  $\|P_{t_i} y\| < \delta_i$  for  $i \in \mathbf{N}$ ,

where  $P_j$  is the canonical projection from  $Y$  onto  $E_j$ . Let  $\tilde{F}_n = \bigoplus_{i=l(n)+1}^{l(n+1)} F_i$ , and let  $\tilde{P}_j$  be the canonical projection from  $X$  onto  $\tilde{F}_j$ . Since  $(x_A)$  is a weakly null tree, we can pick inductively a branch  $(x_{A_i})$  and an increasing sequence of integers  $1 = k_0 < k_1 < \dots$  such that for any  $i \in \mathbf{N}$ , we have

- (i)  $\|\sum_{j=k_{2i-2}}^{k_{2i-1}-1} \tilde{P}_j x_{A_i} - x_{A_i}\| < \delta_i$ ; and
- (ii)  $\|\sum_{j=k_{2i-2}}^{k_{2i-1}-1} \tilde{P}_j x_{A_t}\| < \delta_{\max\{i,t\}}$  for any  $t \neq i$ .

We prove further that  $(x_{A_i})$  is unconditional. Let  $x = \sum a_i x_{A_i}$ ,  $\|x\| = 1$ . Let  $y \in S_Y$  so that  $Q(y) = x$ . Then  $y$  can be written as  $\sum y_j$ , where  $(y_j)$  satisfies (I)–(IV). Define  $k_{-1} = -1$ , and let  $z_i = \sum_{j=k_{2i-3}+2}^{k_{2i-1}+1} y_j$ . We prove that  $\|Qz_i - a_i x_{A_i}\|$  is small:

$$\begin{aligned} \|Qz_i - a_i x_{A_i}\| &\leq \left\| Q\left(\sum_{j=t_{k_{2i-3}+1}}^{t_{k_{2i-1}+1}} P_j y\right) - \left(\sum_{j=k_{2i-2}}^{k_{2i-1}-1} \tilde{P}_j\right)x \right\| + \left\| z_i - \sum_{j=t_{k_{2i-3}+1}}^{t_{k_{2i-1}+1}} P_j y \right\| \\ &\quad + \left\| a_i x_{A_i} - \left(\sum_{j=k_{2i-2}}^{k_{2i-1}-1} \tilde{P}_j\right)x \right\|. \end{aligned} \tag{2.1}$$

Hence we need to estimate the three terms on the right-hand side of the above inequality. By the construction, for  $i > 1$  we have

$$\begin{aligned} \left\| z_i - \sum_{j=t_{k_{2i-3}+1}}^{t_{k_{2i-1}+1}} P_j y \right\| &< \sum_{j=k_{2i-3}+2}^{k_{2i-1}+1} \left( \left\| \sum_{l=t_{j-1}+1}^{t_j-1} P_l y - y_j \right\| + \|P_{t_{j-1}} y\| \right) + \|P_{t_{k_{2i-1}+1}} y\| \\ &< \sum_{j=k_{2i-3}+2}^{k_{2i-1}+1} \delta_j + \sum_{j=k_{2i-3}+2}^{k_{2i-1}+2} \delta_{j-1} < \delta_{k_{2i-3}+1} + \delta_{k_{2i-3}} < \delta_i. \end{aligned} \tag{2.2}$$

By direct calculation, for  $i = 1$  we have

$$\left\| z_1 - \sum_{j=1}^{t_{k_1+1}} P_j y \right\| < 2\delta_1. \tag{2.3}$$

This gives an estimate of the second term. For the third term, we have

$$\begin{aligned} \left\| a_i x_{A_i} - \left( \sum_{j=k_{2i-2}}^{k_{2i-1}-1} \tilde{P}_j \right) x \right\| &< \left\| \left( \sum_{j=k_{2i-2}}^{k_{2i-1}-1} \tilde{P}_j \right) (a_i x_{A_i} - x) \right\| + \left\| a_i x_{A_i} - \left( \sum_{j=k_{2i-2}}^{k_{2i-1}-1} \tilde{P}_j \right) x_{A_i} \right\| \\ &< 2 \left( k_{2i-2} \delta_{k_{2i-2}} + \sum_{j \geq k_{2i-1}} \delta_j \right) + 2\delta_i \\ &< 2(\delta_{k_{2i-2}-1} + \delta_{k_{2i-1}-1}) + 2\delta_i < 4\delta_i. \end{aligned} \tag{2.4}$$

For the first term, let  $Q_j$  be the canonical projection from  $X$  onto  $F_j$ , and let  $J_1 = [t_{k_{2i-3}+1}, t_{k_{2i-1}+1}]$ ,  $J_2 = [l_{k_{2i-2}} + 1, l_{k_{2i-1}}]$ , and  $J'_1 = (t_{k_{2i-3}+1}, t_{k_{2i-1}+1})$ . Then we have

$$\begin{aligned} &\left\| Q \left( \sum_{j \in J_1} P_j y \right) - \left( \sum_{j \in J_2} Q_j \right) Q y \right\| \\ &\leq \left\| Q \left( \sum_{j \in J_1} P_j y \right) - \left( \sum_{j \in J_1} Q_j \right) Q y \right\| + \left\| \left( \sum_{j \in J_1} Q_j \right) Q y - \left( \sum_{j \in J_2} Q_j \right) Q y \right\| \\ &= \left\| Q \left( \sum_{j \in J_1} P_j y \right) - \left( \sum_{j \in J_1} Q_j \right) Q y \right\| + \left\| \sum_{j \in J_1 - J_2} Q_j \left( \sum a_i x_{A_i} \right) \right\| \\ &< \left\| Q \left( \sum_{j \in J_1} P_j y \right) - \left( \sum_{j \in J_1} Q_j \right) Q y \right\| + 4\delta_i \\ &\leq \left\| \left( \sum_{j \notin J_1} Q_j \right) Q \left( \sum_{j \in J_1} P_j y \right) \right\| + \left\| \left( \sum_{j \in J_1} Q_j \right) Q \left( \sum_{j \notin J_1} P_j y \right) \right\| + 4\delta_i \\ &< \left\| \left( \sum_{j \notin J_1} Q_j \right) Q \left( \sum_{j \in J'_1} P_j y \right) \right\| + \left\| \left( \sum_{j \in J'_1} Q_j \right) Q \left( \sum_{j \notin J_1} P_j y \right) \right\| + 6\delta_i \\ &< 2\lambda\delta_i + 2\lambda\delta_i + 6\delta_i = (4\lambda + 6)\delta_i. \end{aligned} \tag{2.5}$$

From inequalities (2.2)–(2.5), we conclude that

$$\|Qz_i - a_i x_{A_i}\| < (4\lambda + 12)\delta_i.$$

Let  $(\epsilon_i) \subset \{-1, 1\}^{\mathbb{N}}$ . Let  $I \subset \mathbb{N}$  be the set of indices  $i \in \mathbb{N}$  for which  $\|y_i\| < \delta_i$ , and let  $I_i = [k_{2i-3} + 2, k_{2i-1} + 1]$ . So  $z_i = \sum_{j \in I_i} y_j$ . Let  $z'_i = \sum_{j \in I_i - 1} y_j$ . It is easy to verify that  $\|z_i - z'_i\| < \delta_i$ . Hence  $\|Qz'_i - a_i x_{A_i}\| < (4\lambda + 13)\delta_i$ . Now, by (II), we know that  $(z'_i)$  is a  $\delta$ -skipped block sequence. Hence  $(z'_i)$  is unconditional. So we have

$$\begin{aligned} \left\| \sum \epsilon_i a_i x_{A_i} \right\| &\leq \left\| Q \left( \sum \epsilon_i z'_i \right) \right\| + (4\lambda + 13)\epsilon \leq C \left\| \sum z'_i \right\| + (4\lambda + 13)\epsilon \\ &< C \left( \left\| \sum z_i \right\| + \sum \delta_i \right) + (4\lambda + 13)\epsilon \leq C + (C + 4\lambda + 13)\epsilon. \end{aligned}$$

This shows that  $(x_{A_i})$  is an unconditional sequence. □

*Remark 2.9*

If the original space  $Y$  has the  $(1 + \epsilon)$ -UTP for any  $\epsilon > 0$ , then any quotient of  $Y$  has the  $(1 + \epsilon)$ -UTP for any  $\epsilon > 0$ .

The following is an elementary lemma, which is used later. We omit the standard proof.

## LEMMA 2.10

Let  $X$  be a Banach space, and let  $X_1, X_2$  be two closed subspaces of  $X$ . If  $X_1 \cap X_2 = \{0\}$  and  $X_1 + X_2$  is closed, then  $X$  embeds into  $X/X_1 \oplus X/X_2$ .

In [7, Theorem 4.4], Johnson and Rosenthal proved that any separable Banach space  $X$  admits a subspace  $Y$  so that both  $Y$  and  $X/Y$  have an FDD. The proof uses Markuschevich bases; a Markuschevich basis for a separable Banach space  $X$  is a biorthogonal system  $\{x_n, x_n^*\}_{n \in \mathbf{N}}$  for which the span of the  $x_n$ 's is dense in  $X$  and the  $x_n^*$ 's separate the points of  $X$ . By [11, Theorem 1.f.4], every separable Banach space  $X$  has a Markuschevich basis  $\{x_n, x_n^*\}_{n \in \mathbf{N}}$  so that  $[x_n^*]$  contains any designated separable subspace of  $X^*$ . The following lemma is a stronger form of the result of Johnson and Rosenthal, which follows from the original proof. For the convenience of the reader, we give a sketch of the proof. We use  $[x_i]_{i \in I}$  to denote the closed linear span of  $(x_i)_{i \in I}$ .

## LEMMA 2.11

Let  $X$  be a separable Banach space. Then there exists a subspace  $Y$  with FDD  $(E_n)$  such that for any blocking  $(F_n)$  of  $(E_n)$  and for any sequence  $(n_k) \subset \mathbf{N}$ ,  $X/\overline{\text{span}\{(F_{n_k})_{k=1}^\infty\}}$  admits an FDD  $(G_n)$ . Moreover, if  $X^*$  is separable,  $(E_n)$  and  $(G_n)$  can be chosen to be shrinking.

*Proof*

Let  $\{x_i, x_i^*\}$  be a Markuschevich basis for  $X$  so that  $[x_i^*]$  is a norm-determining subspace of  $X^*$  and even  $[x_i^*] = X^*$  if  $X^*$  is separable. Then we can choose inductively finite sets  $\sigma_1 \subset \sigma_2 \subset \dots$  and  $\eta_1 \subset \eta_2 \subset \dots$  so that  $\sigma = \bigcup_{n=1}^\infty \sigma_n$  and  $\eta = \bigcup_{n=1}^\infty \eta_n$  are complementary infinite subsets of the positive integers and for  $n = 1, 2, \dots$ ,

- (i) if  $x^* \in [x_i^*]_{i \in \eta_n}$ , there is an  $x \in [x_i]_{i \in \eta_n \cup \sigma_{n+1}}$  such that  $\|x\| = 1$  and  $|x^*(x)| > (1 - 1/(n+1))\|x^*\|$ ;
- (ii) if  $x \in [x_i]_{i \in \sigma_n}$ , there is an  $x^* \in [x_i^*]_{i \in \sigma_n \cup \eta_n}$  such that  $\|x^*\| = 1$  and  $|x^*(x)| > (1 - 1/(n+1))\|x\|$ .

Once we have this, by [7, proof of Theorem 4] we have it that  $[x_i]_{i \in \sigma}^\perp$  is the  $w^*$ -closure of  $[x_i^*]_{i \in \eta}$ . Put  $Y = [x_i^*]_{i \in \eta}^\perp = [x_i]_{i \in \sigma}$ . By the analogue of [7, Proposition 2.1(a)], we deduce that  $X/Y$  has an FDD and that  $([x_i]_{i \in \sigma_n})_{n=1}^\infty$  forms an FDD for  $Y$ . In order to prove Lemma 2.11, it is enough to prove that for any blocking  $(\Sigma_n)$  of  $(\sigma_n)$  or any subsequence  $(\sigma_{n_k})$  of  $(\sigma_n)$  (this, of course, needs the redefining of  $(\eta_n)$ ),

(i) and (ii) still hold. But this is more or less obvious because if  $\Sigma_n = \bigcup_{i=k_{n-1}+1}^{k_n} \sigma_i$ , then we define  $\Delta_n = \bigcup_{i=k_{n-1}+1}^{k_n} \eta_i$  and it is easy to check that  $\{\Sigma_n, \Delta_n\}$  satisfy (i) and (ii). For a subsequence  $(\sigma_{n_k})$ , if we let  $\Sigma_k = \sigma_{n_k}$  and define  $\Delta_k = \bigcup_{i=n_k}^{n_{k+1}-1} \eta_i$ , then  $\{\Sigma_n, \Delta_n\}$  satisfy (i) and (ii). The rest is exactly the same as in [7, proof of Theorem 4.4].  $\square$

The next lemma shows that for a reflexive space with a USB FDD, its dual also has a USB FDD.

LEMMA 2.12

Let  $X$  be a reflexive Banach space with a USB FDD  $(E_n)$ . Then there is a blocking  $(F_n)$  of  $(E_n)$  such that  $(F_n^*)$  is a USB FDD for  $X^*$ .

*Proof*

Without loss of generality, we assume that  $(E_n)$  is monotone. Let  $(\delta_i)$  be a sequence of positive numbers decreasing fast to zero. By the “killing the overlap” technique, we get a blocking  $(F_n)$  of  $(E_n)$  with  $F_n = \sum_{i=k_{n-1}+1}^{k_n} E_i$  so that given any  $x = \sum x_i$  with  $x_i \in E_i, \|x\| = 1$ , there is an increasing sequence  $(t_n)$  with  $k_{n-1} < t_n < k_n$  such that  $\|x_{t_i}\| < \delta_i$ , where  $0 = k_0 < k_1 < \dots$ . Let  $(F_n^*)$  be the dual FDD of  $(F_n)$ , and let  $(x_i^*)$  be a normalized skipped block sequence with respect to  $(F_n^*)$  so that  $x_i^* \in \bigoplus_{j=m_{i-1}+1}^{m_i-1} F_j^*$ , where  $0 = m_0 < m_1 < \dots$ . Let  $x^* = \sum a_i x_i^*$  with  $\|x^*\| = 1$ . Let  $x = \sum x_i$  be a norming functional of  $x^*$  with  $x_i \in E_i$ . By the definition of  $(F_n)$ , we get an increasing sequence  $(t_i)$  with  $k_{i-1} < t_i < k_i$  so that  $\|x_{t_i}\| < \delta_i$ . We define  $y_1 = \sum_{j=1}^{t_{m_1}-1} x_j$  and  $y_i = \sum_{j=t_{m_{i-1}}+1}^{t_{m_i}-1} x_j$  for  $i > 1$ . Let  $y = \sum y_i$ . So by the triangle inequality,

$$\|x - y\| \leq \left\| \sum x_{t_{m_i}} \right\| \leq \sum \|x_{t_{m_i}}\| < \sum \delta_{t_{m_i}}.$$

Let  $(\epsilon_i) \subset \{-1, 1\}^{\mathbb{N}}$ , and let  $\tilde{x}^* = \sum \epsilon_i a_i x_i^*$ . We estimate  $\tilde{x}^*(\sum \epsilon_i y_i)$ :

$$\begin{aligned} \left| \tilde{x}^*\left(\sum \epsilon_i y_i\right) \right| &= \left| \sum \epsilon_i a_i x_i^*\left(\sum \epsilon_i y_i\right) \right| = \left| \sum a_i x_i^*\left(\sum y_i\right) \right| \\ &= |x^*(y)| \geq 1 - \sum \delta_{t_{m_i}}. \end{aligned}$$

Since  $(y_i)$  is a skipped block sequence with respect to  $(E_i)$ ,  $(y_i)$  is unconditional. Hence

$$\left\| \sum \epsilon_i y_i \right\| \leq C \left\| \sum y_i \right\| < C \left(1 + \sum \delta_{m_i}\right),$$

where  $C$  is the unconditional constant associated with the USB FDD  $(E_n)$ . If we let  $\sum \delta_i < \epsilon/2$ , then we conclude that

$$\|\tilde{x}^*\| > \frac{1 - \epsilon}{C(1 + \epsilon)}.$$

Therefore  $(x_i^*)$  is unconditional with unconditional constant less than  $(1 + 3\epsilon)C$  if  $\epsilon$  is sufficiently small. Hence  $(F_n^*)$  is a USB FDD.  $\square$

#### THEOREM 2.13

*Let  $X$  be a separable reflexive Banach space. Then the following are equivalent.*

- (a)  $X$  has the UTP.
- (b)  $X$  embeds into a reflexive Banach space with a USB FDD.
- (c)  $X^*$  has the UTP.

#### *Proof*

It is obvious that (b) implies (a). If we can prove that (a) implies (b) and that  $X$  satisfies (b), then by Lemma 2.12,  $X^*$  is a quotient of a reflexive space with a USB FDD. So, by Theorem 2.8,  $X^*$  has the UTP. Hence we need only show that (a) implies (b). Let  $X_1$  be a subspace of  $X$  with an FDD  $(E_n)$  given by Lemma 2.11. By [13, Proposition 2.4], we get a blocking  $(F_n)$  of  $(E_n)$  so that  $(F_n)$  is a USB FDD. Let  $Y_1 = [F_{4n}]$ , and let  $Y_2 = [F_{4n+2}]$ . Then  $(F_{4n})$  and  $(F_{4n+2})$  form UFDDs for  $Y_1$  and  $Y_2$ . By Lemma 2.11,  $X/Y_i$  has an FDD. Since  $X$  has the UTP, by Theorem 2.8 we know that  $X/Y_i$  has the UTP. By using [13, Proposition 2.4] again, we know that  $X/Y_i$  has a USB FDD. Noticing that  $Y_1 \cap Y_2 = \{0\}$  and that  $Y_1 + Y_2$  is closed, by Lemma 2.10 we have that  $X$  embeds into  $X/Y_1 \oplus X/Y_2$ . Hence  $X$  embeds into a reflexive space with a USB FDD.  $\square$

#### COROLLARY 2.14

*Let  $X$  be a separable reflexive Banach space with the UTP. Then  $X$  embeds into a reflexive Banach space with an unconditional basis.*

#### *Proof*

By Theorem 2.13,  $X$  embeds into a reflexive space  $Y$  with a USB FDD  $(E_n)$ . We prove that  $Y$  embeds into a reflexive space with a UFDD. Then, as mentioned in the introduction,  $Y$  embeds into a reflexive space with an unconditional basis, and so  $X$  does, too.

By Lemma 2.12, there is a blocking  $(F_n)$  of  $(E_n)$  such that  $(F_n^*)$  is a USB FDD for  $Y^*$ . Now, let  $Y_1 = \bigoplus F_{4n}$ , and let  $Y_2 = \bigoplus F_{4n+2}$ . Then we have  $Y_1 \cap Y_2 = \{0\}$ , and  $Y_1 + Y_2$  is closed because  $(F_{2n})$ , being a skipped blocking of  $(E_n)$ , is unconditional. By Lemma 2.10,  $Y$  embeds into  $Y/Y_1 \oplus Y/Y_2$ . Since  $(Y/Y_i)^*$  is isomorphic to  $Y_i^\perp$ , it

is enough to prove that  $Y_i^\perp$  has a UFDD. Let  $G_n^* = F_{4n-3}^* \oplus F_{4n-2}^* \oplus F_{4n-1}^*$ . It is easy to see that  $(G_n^*)$  forms an FDD for  $Y_1^\perp$ . Noticing that  $(G_n)$  is a skipped blocking of  $(F_n^*)$ , we conclude that  $(G_n)$  is unconditional. Similarly, we can show that  $Y_2^\perp$  admits a UFDD. This finishes the proof.  $\square$

**COROLLARY 2.15**

*Let  $X$  be a quotient of a reflexive Banach space with a UFDD. Then  $X$  embeds into a reflexive Banach space with an unconditional basis.*

*Proof*

Combine Theorem 2.8 and Corollary 2.14.  $\square$

We mention again that in 1974, Davis, Figiel, Johnson, and Pełczyński proved in [1] that a reflexive Banach space  $X$  that embeds into a Banach space with a shrinking unconditional basis embeds into a reflexive space  $X$  with an unconditional basis. The next year, Figiel, Johnson, and Tzafriri [4] got a stronger result by removing the shrinkingness of the unconditional basis in the hypothesis. Our next corollary gives a parallel result for quotients.

**COROLLARY 2.16**

*Let  $X$  be a separable reflexive Banach space. If  $X$  is a quotient of a Banach space with a shrinking unconditional basis, then  $X$  is isomorphic to a quotient of a reflexive Banach space with an unconditional basis.*

*Proof*

Since  $X$  is a quotient of a Banach space with a shrinking unconditional basis,  $X^*$  is a subspace of a Banach space with an unconditional basis. Hence by [4, Theorem 3.1],  $X^*$  is isomorphic to a subspace of a reflexive Banach space with an unconditional basis. Therefore,  $X$  is isomorphic to a quotient of a reflexive Banach space with an unconditional basis.  $\square$

**Remark 2.17**

Corollary 2.16 is different from the result of [4] in that the shrinkingness in our result cannot be removed. The reason is more or less obvious since every separable Banach space is a quotient of  $\ell_1$ , which has an unconditional basis.

Gluing Theorem 2.13 and Corollaries 2.14, 2.15, and 2.16 together, we have the following long list of equivalences.

**THEOREM 2.18**

*Let  $X$  be a separable reflexive Banach space. Then the following are equivalent.*

- (a)  $X$  has the UTP.
- (b)  $X$  is isomorphic to a subspace of a Banach space with an unconditional basis.
- (c)  $X$  is isomorphic to a subspace of a reflexive space with an unconditional basis.
- (d)  $X$  is isomorphic to a quotient of a Banach space with a shrinking unconditional basis.
- (e)  $X$  is isomorphic to a quotient of a reflexive space with an unconditional basis.
- (f)  $X$  is isomorphic to a subspace of a quotient of a reflexive space with an unconditional basis.
- (g)  $X$  is isomorphic to a subspace of a reflexive quotient of a Banach space with a shrinking unconditional basis.
- (h)  $X$  is isomorphic to a quotient of a subspace of a reflexive space with an unconditional basis.
- (i)  $X$  is isomorphic to a quotient of a reflexive subspace of a Banach space with a shrinking unconditional basis.

**3. Example**

In this section, we give an example of a reflexive Banach space for which there exists a  $C > 0$  such that every normalized weakly null sequence admits a  $C$ -unconditional subsequence, while for any  $D > 0$  there is a normalized weakly null tree such that every branch is not  $D$ -unconditional. The construction is an analogue of Odell and Schlumprecht’s example (see [12, Example 4.2]).

We first construct an infinite sequence of reflexive Banach spaces  $X_n$ . Each  $X_n$  is infinite-dimensional and has the property that for  $\epsilon > 0$ , every normalized weakly null sequence has a  $(1 + \epsilon)$ -unconditional basic subsequence, while there is a normalized weakly null tree for which every branch is at least  $C_n$ -unconditional and  $C_n$  goes to infinity when  $n$  goes to infinity. Then the  $\ell_2$ -sum of  $X_n$ ’s is a reflexive Banach space with the desired property.

Let  $[\mathbf{N}]^{\leq n}$  be the set of all subsets of the positive integers with cardinality less than or equal to  $n$ . Let  $c_{00}([\mathbf{N}]^{\leq n})$  be the space of sequences with finite support indexed by  $[\mathbf{N}]^{\leq n}$ , and denote its canonical basis by  $(e_A)_{A \in [\mathbf{N}]^{\leq n}}$ . Let  $(h_i)$  be any normalized conditional basic sequence that satisfies a block lower  $\ell_2$ -estimate with constant 1, for example, the boundedly complete basis of James’s space (see [2, Problem 6.41]). Let  $\sum a_A e_A$  be an element of  $c_{00}([\mathbf{N}]^{\leq n})$ . Let  $(\beta_k)_{k=1}^m$  be disjoint segments. By “a segment in  $[\mathbf{N}]^{\leq n}$ ,” we mean a sequence  $(A_i)_{i=1}^k \in [\mathbf{N}]^{\leq n}$  with  $A_1 = \{n_1, n_2, \dots, n_l\}$ ,  $A_2 = \{n_1, n_2, \dots, n_l, n_{l+1}\}$ ,  $\dots$ ,  $A_k = \{n_1, n_2, \dots, n_l, \dots, n_{l+k-1}\}$  for some  $n_1 < n_2 < \dots < n_{l+k-1}$ . Let  $\beta_k = \{A_{1,k}, A_{2,k}, \dots, A_{j_k,k}\}$  with  $A_{i,k} < A_{i+1,k}$  under the tree order in  $[\mathbf{N}]^{\leq n}$ . Now, we define  $X_n$  to be the completion of  $c_{00}([\mathbf{N}]^{\leq n})$  under the norm

$$\left\| \sum a_A e_A \right\|_{X_n} = \sup \left\{ \left( \sum_{k=1}^m \left( \left\| \sum_{A_{i,k} \in \beta_k} a_{A_{i,k}} h_i \right\| \right)^2 \right)^{1/2} : (\beta_k)_{k=1}^m \text{ are disjoint segments} \right\}.$$

Let  $X = \left(\sum X_n\right)_2$ . Let  $C_M$  be the unconditional constant of  $(h_i)_{i=1}^M$ . It is clear that  $C_M$  tends to infinity when  $M$  goes to infinity. The normalized weakly null tree  $(e_A)_{A \in [\mathbf{N}]^{\leq M}}$  in  $X_M$  has the property that every branch of it is 1-equivalent to  $(h_i)_{i=1}^M$  since  $(h_i)$  has a block lower  $\ell_2$ -estimate with constant 1. So what is remaining is to verify that for every  $\epsilon > 0$ , every normalized weakly null sequence in  $X$  has a  $(1 + \epsilon)$ -unconditional basic subsequence. Actually, we prove that there is a subsequence which is  $(1 + \epsilon)$ -equivalent to the unit vector basis of  $\ell_2$ . By a gliding-hump argument, it is not hard to verify the following fact.

*Fact*

Let  $(Y_k)$  be a sequence of reflexive Banach spaces, and let  $Y = \left(\sum Y_k\right)_{\ell_2}$ . If for every  $\epsilon > 0$ ,  $k \in \mathbf{N}$ , every normalized weakly null sequence in  $Y_k$  has a subsequence that is  $(1 + \epsilon)$ -equivalent to the unit vector basis of  $\ell_2$ , then for every  $\epsilon > 0$ , every normalized weakly null sequence in  $Y$  has a subsequence that is  $(1 + \epsilon)$ -equivalent to the unit vector basis of  $\ell_2$ .

Considering this fact, it is enough to show that for every  $\epsilon > 0$ ,  $k \in \mathbf{N}$ , every normalized weakly null sequence in  $X_k$  has a subsequence that is  $(1 + \epsilon)$ -equivalent to the unit vector basis of  $\ell_2$ . We prove this by induction.

For  $k = 1$ ,  $X_1$  is isometric to  $\ell_2$ , so the conclusion is obvious.

Assume that the conclusion is true for  $X_k$ . By the definition of  $X_{k+1}$ ,  $X_{k+1}$  is isometric to  $\left(\sum(\mathbf{R} \oplus X_k)\right)_{\ell_2}$  (where  $\mathbf{R} \oplus X_k$  has some norm so that  $\{0\} \oplus X_k$  is isometric to  $X_k$ ). Hence by hypothesis and the fact mentioned above, it is easy to see that the conclusion is true in  $X_{k+1}$ . This finishes the proof.

*Remark 3.1*

The proof of the corresponding induction step in [12, Example 4.2] is more complicated than the very simple induction argument in the previous paragraph. Schlumprecht realized after [12] was published that the induction could be done this simply (see [16]), and his argument works in our context.

*Acknowledgments.* The authors thank the referees for useful corrections, especially for pointing out the imprecision in the initial construction of the example in Section 3. This article is based in part on the doctoral dissertation of Zheng, which is being prepared at Texas A&M University under Johnson's direction.

**References**

- [1] W. J. DAVIS, T. FIGIEL, W. B. JOHNSON, and A. PEŁCZYŃSKI, *Factoring weakly compact operators*, J. Functional Analysis **17** (1974), 311–327. MR 0355536

- [2] M. FABIAN, P. HABALA, P. HÁJEK, V. MONTESINOS SANTALUCÍA, J. PELANT, and V. ZIZLER, *Functional Analysis and Infinite-Dimensional Geometry*, CMS Books Math./Ouvrages Math. SMC **8**, Springer, New York, 2001. MR 1831176
- [3] M. FEDER, *On subspaces of spaces with an unconditional basis and spaces of operators*, Illinois J. Math. **24** (1980), 196–205. MR 0575060
- [4] T. FIGIEL, W. B. JOHNSON, and L. TZAFRIRI, *On Banach lattices and spaces having local unconditional structure, with applications to Lorentz function spaces*, J. Approximation Theory **13** (1975), 395–412. MR 0367624
- [5] R. HAYDON, E. ODELL, and T. SCHLUMPRECHT, *Small subspaces of  $L_p$* , preprint, arXiv:0711.3919v1 [math.FA]
- [6] W. B. JOHNSON, *On quotients of  $L_p$  which are quotients of  $\ell_p$* , Compositio Math. **34** (1977), 69–89. MR 0454595
- [7] W. B. JOHNSON and H. P. ROSENTHAL, *On  $\omega^*$ -basic sequences and their applications to the study of Banach spaces*, Studia Math. **43** (1972), 77–92. MR 0310598
- [8] W. B. JOHNSON and A. SZANKOWSKI, *Complementably universal Banach spaces*, Studia Math. **58** (1976), 91–97. MR 0425582
- [9] W. B. JOHNSON and M. ZIPPIN, “On subspaces of quotients of  $(\sum G_n)_{i_p}$  and  $(\sum G_n)_{c_0}$ ” in *Proceedings of the International Symposium on Partial Differential Equations and the Geometry of Normal Linear Spaces (Jerusalem, 1972)*, Israel J. Math. **13** (1972), 311–316. MR 0331023
- [10] ———, *Subspaces and quotient spaces of  $(\sum G_n)_{i_p}$  and  $(\sum G_n)_{c_0}$* , Israel J. Math. **17** (1974), 50–55. MR 0358296
- [11] J. LINDENSTRAUSS and L. TZAFRIRI, *Classical Banach Spaces, I: Sequence Spaces*, Ergeb. Math. Grenzgeb. **92**, Springer, Berlin, 1977. MR 0500056
- [12] E. ODELL and T. SCHLUMPRECHT, *Trees and branches in Banach spaces*, Trans. Amer. Math. Soc. **354**, no. 10 (2002), 4085–4108. MR 1926866
- [13] ———, *A universal reflexive space for the class of uniformly convex Banach spaces*, Math. Ann. **335** (2006), 901–916. MR 2232021
- [14] E. ODELL, T. SCHLUMPRECHT, and A. ZSÁK, *On the structure of asymptotic  $\ell_p$  spaces*, to appear in Q. J. Math.
- [15] A. PEŁCZYŃSKI and P. WOJTASZCZYK, *Banach spaces with finite dimensional expansions of identity and universal bases of finite dimensional subspaces*, Studia Math. **40** (1971), 91–108. MR 0313765
- [16] T. SCHLUMPRECHT, private communication, 2006.
- [17] M. ZIPPIN, *Banach spaces with separable duals*, Trans. Amer. Math. Soc. **310**, no. 1 (1988), 371–379. MR 0965758

*Johnson*

Department of Mathematics, Texas A&M University, College Station, Texas 77843, USA;  
johnson@math.tamu.edu

*Zheng*

Department of Mathematics, University of Texas at Austin, Austin, Texas 78712, USA;  
btzheng@math.utexas.edu